

Quadratic Forms and Space-Time Block Codes from Generalized Quaternion and Biquaternion Algebras

Thomas Unger and Nadya Markin

Abstract—In the context of space-time block codes (STBCs), the theory of generalized quaternion and biquaternion algebras (i.e., tensor products of two quaternion algebras) over arbitrary base fields is presented, as well as quadratic form theoretic criteria to check if such algebras are division algebras. For base fields relevant to STBCs, these criteria are exploited, via Springer’s theorem, to construct several explicit infinite families of (bi-)quaternion division algebras. These can be used to obtain 2×2 and 4×4 STBCs via the left regular representation.

Index Terms—Space-time block codes, division algebras, quadratic forms.

I. INTRODUCTION

EVER since Alamouti’s work [1] a decade ago, division algebras – both commutative (i.e., fields) and noncommutative – have played an increasingly important role in the construction of space-time block codes (STBCs) for wireless communication.

Precisely how this is done is explained very thoroughly in [2]. From a mathematical point of view, one wants to construct, for a given integer n , a set $S \subset M_n(F)$ of $n \times n$ -matrices over some field F such that the difference of any two elements of S has nonzero determinant.

The field F is usually an extension of the field $\mathbb{Q}$ of rational numbers, containing an appropriate signal constellation. The set $S \subset M_n(F)$ can be obtained as follows: choose some division algebra D of dimension n over F (i.e., D is simultaneously an F -vector space and a ring) and consider the left regular representation

$$\lambda : D \rightarrow \text{End}_F(D), \quad a \mapsto \lambda_a,$$

where λ_a is left multiplication by a , $\lambda_a(x) = ax$ for all $x \in D$. Here $\text{End}_F(D)$ is the algebra of F -linear transformations on D . After a choice of F -basis for D , $\text{End}_F(D)$ can be identified with the algebra $M_n(F)$. In this way we get an embedding $\lambda : D \hookrightarrow M_n(F)$. Since all nonzero elements of D are invertible, all nonzero matrices in $\lambda(D)$ will have nonzero determinant. The set S can then be chosen to be (a subset of) $\lambda(D)$. The difference of any two elements of S will thus have nonzero determinant.

There are some caveats here: for coding purposes it is often more convenient to consider D as a vector space over some subfield K , maximal with respect to inclusion ($F \subset K \subset D$). When D is commutative (in other words a field), the above

construction will always work with F replaced by K . However, when D is noncommutative, one should consider D as a *right* K -vector space, i.e., elements of D should be multiplied on the right by scalars from K . It should be taken into account that in both cases $\dim_F D = (\dim_K D)^2$, resulting in K -matrices of smaller size.

We are only interested in noncommutative division algebras in this paper. In the STBC literature two large classes of algebras have been studied: cyclic algebras (e.g. in [2]) and crossed-product algebras (e.g. in [3] and [4]). The latter are characterized by requiring that K is a strictly maximal subfield of D which is a Galois extension of F . Cyclic algebras are crossed-product algebras with *cyclic* Galois group $\text{Gal}(K/F)$. We refer to [5] for a historical survey of such algebras.

Determining when precisely these algebras are *division algebras* can be a nontrivial problem. In this paper we discuss two classes of algebras, namely (generalized) quaternion and biquaternion algebras. They have characterizations in terms of *quadratic forms*, a fact that has not fully been exploited yet in the coding theory literature. There are particularly nice quadratic form-theoretic criteria to determine when they are division algebras.

Although many examples of quaternion algebras have appeared in the literature (first and foremost the Alamouti code of course), they often do so disguised as cyclic algebras (of which they are special cases), which hides their quadratic form-theoretic characterization.

Biquaternion algebras are tensor products of two quaternion algebras (in the older literature, the term also refers to Hamilton’s quaternions considered over the field of complex numbers). They are crossed-product algebras with Galois group the Klein 4-group (but the converse inclusion is not true), and are thus special cases of the algebras in [4]. Biquaternion algebras can also be characterized by a quadratic form. In particular it is relatively straightforward to determine when such an algebra is a division algebra (at least over base fields that are relevant for STBCs). It is also easy to produce biquaternion division algebras as tensor products of quaternion division algebras, addressing an issue raised in [3, p. 3926].

Quaternion division algebras can be used to construct 2×2 STBCs and biquaternion division algebras to construct 4×4 STBCs.

This paper is organized as follows: in Section II we present a brief introduction to quadratic forms. In Sections III and IV we respectively present the theory of quaternion and biquaternion algebras over arbitrary fields. We also give general quadratic form theoretic criteria for such algebras to be division algebras. In Section V we obtain matrix representations of quaternion

T. Unger and N. Markin are with the School of Mathematical Sciences, University College Dublin, Belfield, Dublin 4, Ireland (e-mail: thomas.unger@ucd.ie and nadyaomarkin@gmail.com).

Version of June 14, 2010

and biquaternion algebras over arbitrary fields by means of the left regular representation.

Then we shift our attention to fields that are actually used in space-time block coding: number fields and their transcendental extensions. Such fields can be embedded into bigger fields that are complete with respect to a discrete valuation, or CDV fields for short. For CDV fields the quadratic form theoretic criterium for a (bi-)quaternion algebra to be division can be tested in a finite number of steps by means of Springer's theorem (Section VI).

In the final two sections we give examples of infinite families of non-isomorphic quaternion and biquaternion algebras over suitable fields, exploiting Springer's theorem. Together with the matrix representations from Section V these can be used for the explicit construction of 2×2 and 4×4 STBCs.

The Appendices contain preliminaries from discrete valuation theory and number theory.

II. QUADRATIC FORMS

For the convenience of the reader we recall some basic facts from quadratic form theory. We refer to [6] for the details. Let F be a field and write $F^\times := F \setminus \{0\}$. We make the technical assumption that the characteristic of F is different from 2 (in other words that 2 is invertible in F). The fields considered in this paper are all extensions of the rational numbers $\mathbb{Q}$ and thus will satisfy this condition (but the finite field $\mathbb{F}_2$ does not, for example).

Let V be a finite dimensional F -vector space.

Definition 2.1: A *quadratic form* on V (over F) is a map $\varphi : V \rightarrow F$ satisfying the following two properties:

- (i) $\varphi(\alpha \mathbf{v}) = \alpha^2 \varphi(\mathbf{v})$ for all $\mathbf{v} \in V$ and all $\alpha \in F$;
- (ii) the map $b_\varphi : V \times V \rightarrow F$ defined by

$$b_\varphi(\mathbf{v}, \mathbf{w}) := \varphi(\mathbf{v} + \mathbf{w}) - \varphi(\mathbf{v}) - \varphi(\mathbf{w})$$

for all $\mathbf{v}, \mathbf{w} \in V$ is bilinear (and automatically symmetric).

Concretely, if we identify V with F^n , where $n = \dim_F V$, then φ is just a homogeneous polynomial of degree two: for $1 \leq i, j \leq n$ there exist scalars $a_{ij} \in F$ such that

$$\varphi(\mathbf{v}) = \varphi(v_1, \dots, v_n) = \sum_{i,j=1}^n a_{ij} v_i v_j, \quad (1)$$

for all $\mathbf{v} = (v_1, \dots, v_n) \in F^n$.

If we let $\mathbf{A} = (a_{ij})_{1 \leq i,j \leq n} \in M_n(F)$ we can write (1) in matrix form as

$$\varphi(\mathbf{v}) = \mathbf{v} \mathbf{A} \mathbf{v}^t,$$

where t denotes transposition.

If ψ is another quadratic form on V , with matrix $\mathbf{B}$ say, then we call φ and ψ *isometric* if the matrices $\mathbf{A}$ and $\mathbf{B}$ are congruent, i.e., if there exists an invertible matrix $\mathbf{C}$ such that $\mathbf{B} = \mathbf{C} \mathbf{A} \mathbf{C}^t$. In this case we write $\varphi \simeq \psi$.

One can show that a quadratic form φ is always isometric to a quadratic form ψ whose matrix is diagonal,

$$\psi(\mathbf{v}) = \sum_{i=1}^n a_i v_i^2$$

for all $\mathbf{v} = (v_1, \dots, v_n) \in F^n$ and scalars $a_1, \dots, a_n \in F$. It is convenient to employ the short-hand notation

$$\psi = \langle a_1, \dots, a_n \rangle$$

to indicate that the matrix of ψ has diagonal entries $a_1, \dots, a_n$ and zeroes everywhere else. Thus we can write

$$\varphi \simeq \langle a_1, \dots, a_n \rangle.$$

It is customary to consider quadratic forms *up to isometry*. As a consequence the entries in $\langle a_1, \dots, a_n \rangle$ may be permuted and changed by nonzero squares (i.e., any a_i may be replaced by $\alpha^2 a_i$ for any $\alpha \in F^\times$).

The *dimension* of φ is the integer n (i.e., the dimension of the underlying vector space). If φ' is another quadratic form over F , say of dimension m , $\varphi' \simeq \langle b_1, \dots, b_m \rangle$, then the *orthogonal sum* of φ and φ' , defined by

$$\varphi \perp \varphi' := \langle a_1, \dots, a_n, b_1, \dots, b_m \rangle$$

is a quadratic form on F^{n+m} which is uniquely defined up to isometry.

The quadratic forms φ in this paper will always be *nonsingular* in the sense that their *determinant* $\det \varphi := a_1 a_2 \cdots a_n \in F/F^2$ is always nonzero.

A quadratic form φ on V is called *isotropic over F* if there exists a nonzero vector $\mathbf{v} \in V$ such that $\varphi(\mathbf{v}) = 0$ and *anisotropic over F* otherwise.

Note that an anisotropic form over F may become isotropic after scalar extension to a bigger field.

Remark 2.2: Let $\varphi = \langle a_1, \dots, a_n \rangle$. If there are $b_3, b_4, \dots, b_n \in F^\times$ such that $\varphi \simeq \langle 1, -1, b_3, b_4, \dots, b_n \rangle$, then φ is clearly isotropic. The converse is also true.

III. QUATERNION ALGEBRAS

Definition 3.1: Let $a, b \in F^\times$. A (*generalized*) *quaternion algebra* over F is a 4-dimensional F -vector space with F -basis $\mathcal{B} = \{1, \mathbf{i}, \mathbf{j}, \mathbf{k}\}$, whose ring structure is determined by

$$\mathbf{i}^2 = a, \mathbf{j}^2 = b \quad \text{and} \quad \mathbf{ij} = -\mathbf{ji} = \mathbf{k}. \quad (2)$$

We denote this algebra by $(a, b)_F$. Its elements are called *quaternions*.

It follows from the defining relations (2) that $\mathbf{k}^2 = -ab$ and that $\mathbf{i}, \mathbf{j}$ and $\mathbf{k}$ anti-commute. Addition of quaternions and multiplication of a quaternion by a scalar is component wise, while the product of two quaternions reduces to computing sums of scalar multiples of products of the form $\xi \eta$ with $\xi, \eta \in \mathcal{B}$. The defining relations (2) should be used to simplify these products. Note that the product of two quaternions is not commutative.

Example 3.2: Hamilton's quaternions form the algebra $\mathbb{H} = (-1, -1)_{\mathbb{R}}$.

Let $\mathbf{q} = q_0 + q_1 \mathbf{i} + q_2 \mathbf{j} + q_3 \mathbf{k} \in Q := (a, b)_F$ (with $q_0, q_1, q_2, q_3 \in F$). The *conjugate* of $\mathbf{q}$, denoted $\bar{\mathbf{q}}$, is defined by

$$\bar{\mathbf{q}} := q_0 - q_1 \mathbf{i} - q_2 \mathbf{j} - q_3 \mathbf{k}.$$

The *norm map* $N : Q \rightarrow F$ is defined by $N(\mathbf{q}) := \mathbf{q} \bar{\mathbf{q}}$ for all $\mathbf{q} \in Q$. A straightforward computation shows that

$$N(\mathbf{q}) := \mathbf{q} \bar{\mathbf{q}} = \bar{\mathbf{q}} \mathbf{q} = q_0^2 - a q_1^2 - b q_2^2 + a b q_3^2. \quad (3)$$

Thus we may regard N as a quadratic form in four variables q_0, q_1, q_2, q_3 , called the *norm form* of Q . Using the notation from Section II we write

$$N = \langle 1, -a, -b, ab \rangle.$$

Theorem 3.3: The quaternion algebra $Q = (a, b)_F$ is a division algebra if and only if its norm form $N = \langle 1, -a, -b, ab \rangle$ is anisotropic over F .

Proof: $\Rightarrow$ If $N(\mathbf{q}) = \mathbf{q}\bar{\mathbf{q}}$ were equal to zero for some nonzero $\mathbf{q} \in Q$, then Q would contain zero divisors, contradicting the assumption that Q is a division algebra.

$\Leftarrow$ Let $\mathbf{q}$ be any nonzero element of Q . From $\bar{\mathbf{q}}\mathbf{q} = \mathbf{q}\bar{\mathbf{q}} = N(\mathbf{q}) \neq 0$ we deduce that $\mathbf{q}$ is invertible with $\mathbf{q}^{-1} = \bar{\mathbf{q}}/N(\mathbf{q})$. $\blacksquare$

Example 3.4: Hamilton's quaternion algebra $\mathbb{H} = (-1, -1)_{\mathbb{R}}$ is a division algebra since its norm form $\langle 1, 1, 1, 1 \rangle$ is clearly anisotropic over $\mathbb{R}$. The quaternion algebra $(-1, 1)_{\mathbb{R}}$ is not a division algebra since its norm form $\langle 1, 1, -1, -1 \rangle$ is isotropic over $\mathbb{R}$.

Remark 3.5: Let $Q = (a, b)_F$. One can show that if Q is not a division algebra, that it is then necessarily isomorphic to the full matrix algebra $M_2(F)$. This thus happens precisely when the norm form of Q is isotropic.

Remark 3.6: One can show that two quaternion algebras over F are isomorphic as F -algebras if and only if their norm forms are isometric quadratic forms.

IV. BIQUATERNION ALGEBRAS

Definition 4.1: A *biquaternion algebra* over F is a tensor product of two quaternion algebras over F ,

$$(a, b)_F \otimes_F (c, d)_F,$$

where $a, b, c, d \in F^\times$. Its elements are called *biquaternions*.

Let $\mathcal{B}_1 := \{1, \mathbf{i}_1, \mathbf{j}_1, \mathbf{k}_1\}$ and $\mathcal{B}_2 := \{1, \mathbf{i}_2, \mathbf{j}_2, \mathbf{k}_2\}$ be F -bases for $(a, b)_F$ and $(c, d)_F$, respectively. The elements of $\mathcal{B}_1$ are assumed to commute with the elements of $\mathcal{B}_2$. The ordered set

$$\mathcal{B} := \{\xi \otimes \eta \mid \xi \in \mathcal{B}_1, \eta \in \mathcal{B}_2\}$$

is an F -basis for $B := (a, b)_F \otimes_F (c, d)_F$. Thus $\dim_F B = 16$ and we can express a biquaternion $\mathbf{x} \in B$ uniquely in the form

$$\mathbf{x} = \sum_{\xi \in \mathcal{B}_1} \sum_{\eta \in \mathcal{B}_2} x_{\xi, \eta} \xi \otimes \eta,$$

with $x_{\xi, \eta} \in F$. Addition of biquaternions and multiplication of a biquaternion by a scalar is component wise, while the product of two biquaternions reduces to computing sums of scalar multiples of products of the form

$$(\xi \otimes \eta)(\xi' \otimes \eta') = \xi\xi' \otimes \eta\eta',$$

where $\xi, \xi' \in \mathcal{B}_1$ and $\eta, \eta' \in \mathcal{B}_2$. The defining relations (2) should be used to simplify these products. For example,

$$\begin{aligned} (\mathbf{i}_1 \otimes \mathbf{j}_2)(\mathbf{i}_1 \otimes \mathbf{k}_2) &= \mathbf{i}_1^2 \otimes \mathbf{j}_2\mathbf{k}_2 = a \otimes \mathbf{j}_2\mathbf{i}_2\mathbf{j}_2 = a \otimes (-\mathbf{i}_2\mathbf{j}_2^2) \\ &= a \otimes (-d\mathbf{i}_2) = -ad(1 \otimes \mathbf{i}_2). \end{aligned}$$

Note that the product of two biquaternions is not commutative.

Definition 4.2: Let $B := (a, b)_F \otimes_F (c, d)_F$. The six-dimensional quadratic form

$$\varphi = \langle a, b, -ab - c, -d, cd \rangle$$

over F is called the *Albert form* of B .

This form is named after Adrian A. Albert who carried out much of the pioneering work on the structure of algebras in the 1930s.

Theorem 4.3: The biquaternion algebra $B = (a, b)_F \otimes_F (c, d)_F$ is a division algebra if and only if its Albert form $\varphi = \langle a, b, -ab, -c, -d, cd \rangle$ is anisotropic over F .

The proof of this theorem is beyond the scope of this paper.

Remark 4.4: If the Albert form φ of B is isotropic over F , then one can show that B is isomorphic to the full matrix algebra $M_4(F)$ if and only if φ is isometric to $\langle 1, -1, 1, -1, 1, -1 \rangle$ and also that $B \cong M_2(D)$ for some quaternion division algebra D if and only if $\varphi \simeq \langle 1, -1, e, f, g, h \rangle$ with $\langle e, f, g, h \rangle$ anisotropic.

Remark 4.5: One can show that two biquaternion algebras over F are isomorphic as F -algebras if and only if their Albert forms are *similar*, i.e., isometric up to a multiplicative constant from F .

V. MATRIX REPRESENTATIONS

A. The Quaternion Case

Let $Q = (a, b)_F$ be a quaternion division algebra with F -basis $\{1, \mathbf{i}, \mathbf{j}, \mathbf{k}\}$, so that $\mathbf{i}^2 = a$ and $\mathbf{j}^2 = b$. The field $K := F(\sqrt{a}) = F(\mathbf{i})$ is a maximal subfield of Q with Galois group $\text{Gal}(K/F) = \{1, \sigma\}$ where σ (the nontrivial automorphism of K that fixes F) is determined by $\sigma(\sqrt{a}) = -\sqrt{a}$.

Let $\mathbf{x} = q_0 + q_1\mathbf{i} + q_2\mathbf{j} + q_3\mathbf{k} \in Q$ (with $q_0, q_1, q_2, q_3 \in F$). Since

$$\mathbf{x} = q_0 + q_1\mathbf{i} + q_2\mathbf{j} + q_3\mathbf{k} = (q_0 + q_1\mathbf{i}) + \mathbf{j}(q_2 - q_3\mathbf{i}) = x_0 + \mathbf{j}x_1$$

with $x_0 = q_0 + q_1\mathbf{i}$ and $x_1 = q_2 - q_3\mathbf{i}$ in K , we can write

$$Q = F(\mathbf{i}) \oplus \mathbf{j}F(\mathbf{i}) = K \oplus \mathbf{j}K.$$

Hence Q can be considered as a two-dimensional right K -vector space with K -basis $\{1, \mathbf{j}\}$. We thus multiply elements $\mathbf{x}$ from Q with scalars α from K on the *right* to get $\mathbf{x}\alpha$. In computations we may bump into expressions of the form $\alpha\mathbf{x}$ with scalars on the left. They can be brought into the correct form by means of the ‘‘hopping over’’ rule

$$\alpha\mathbf{x} = \mathbf{x}\sigma(\alpha) \quad (4)$$

(which can be verified to be true by direct computation).

Theorem 5.1:

$$Q \cong \left\{ \begin{bmatrix} x_0 & b\sigma(x_1) \\ x_1 & \sigma(x_0) \end{bmatrix} \mid x_0, x_1 \in K \right\}$$

Proof: Let $\lambda : Q \hookrightarrow \text{End}_K(Q)$, $\mathbf{x} \mapsto \lambda_{\mathbf{x}}$ be the left regular representation. We can identify $\text{End}_K(Q)$ with $M_2(K)$ by mapping $\lambda_{\mathbf{x}}$ to its matrix with respect to the K -basis $\{1, \mathbf{j}\}$

of Q , which we will denote by $\mathbf{X}$. Let $\mathbf{x} = x_0 + \mathbf{j}x_1 \in Q$. Since

$$\begin{aligned}\lambda_{\mathbf{x}}(1) &= \mathbf{x} = x_0 + \mathbf{j}x_1, \\ \lambda_{\mathbf{x}}(\mathbf{j}) &= \mathbf{x}\mathbf{j} = (x_0 + \mathbf{j}x_1)\mathbf{j} = x_0\mathbf{j} + \mathbf{j}x_1\mathbf{j} = \mathbf{j}\sigma(x_0) + \mathbf{j}^2\sigma(x_1) \\ &= b\sigma(x_1) + \mathbf{j}\sigma(x_0)\end{aligned}$$

(where we used the “hopping over” rule (4) and the fact that $\mathbf{j}^2 = b$), we obtain

$$\mathbf{X} = \begin{bmatrix} x_0 & b\sigma(x_1) \\ x_1 & \sigma(x_0) \end{bmatrix}.$$

Since the map $\mathbf{x} \mapsto \mathbf{X}$ yields an isomorphism of Q with its image inside $M_2(K)$ we are done. $\blacksquare$

Example 5.2: Letting $F = \mathbb{R}$, $K = \mathbb{C}$ and $Q = (-1, -1)_{\mathbb{R}}$ we obtain the Alamouti code.

More examples will be constructed in Section VII.

Remark 5.3: Note that

$$\det \mathbf{X} = x_0\sigma(x_0) - bx_1\sigma(x_1) = q_0^2 - aq_1^2 - bq_2^2 + abq_3^2 = N(\mathbf{x}),$$


confirming once again that invertible elements of Q (i.e., elements with nonzero norm) correspond to invertible matrices.

Remark 5.4: The field $K = F(\sqrt{a})$ is a so-called *splitting field* for Q in the sense that $Q \otimes_F K \cong M_2(K)$.

B. The Biquaternion Case

Let $B := (a, b)_F \otimes_F (c, d)_F$ be a biquaternion division algebra. Let $\mathcal{B}_1, \mathcal{B}_2$ and $\mathcal{B}$ be as in Section IV. Then $\mathbf{i}_1^2 = a$, $\mathbf{j}_1^2 = b$, $\mathbf{i}_2^2 = c$ and $\mathbf{j}_2^2 = d$. The field $K := F(\sqrt{a}, \sqrt{c})$ is a maximal subfield of B with Galois group $\text{Gal}(K/F) = \{1, \sigma, \tau, \sigma\tau\}$ where σ and τ are determined by $\sigma(\sqrt{a}) = -\sqrt{a}$ and $\tau(\sqrt{c}) = -\sqrt{c}$, respectively, and where $\sigma\tau$ denotes the composition of σ and τ .

The field K is a biquadratic extension of the field F . The following diagram illustrates the relationship between K , $F(\sqrt{a})$, $F(\sqrt{c})$ and F :



A K -basis for B is given by $\{1 \otimes 1, 1 \otimes \mathbf{j}_2, \mathbf{j}_1 \otimes 1, \mathbf{j}_1 \otimes \mathbf{j}_2\}$. One can easily verify that B is a right K -vector space which can be identified with

$$K \oplus \mathbf{j}_1 K \oplus \mathbf{j}_2 K \oplus \mathbf{j}_1 \mathbf{j}_2 K.$$

This time there are several “hopping over” rules for multiplication by scalars on the left, determined by

$$\alpha \mathbf{j}_1 = \mathbf{j}_1 \sigma(\alpha), \quad \alpha \mathbf{j}_2 = \mathbf{j}_2 \tau(\alpha) \quad \text{and} \quad \alpha \mathbf{j}_1 \mathbf{j}_2 = \mathbf{j}_1 \mathbf{j}_2 \sigma\tau(\alpha) \quad (5)$$

for all $\alpha \in K$.

Theorem 5.5:

$$B \cong \left\{ \begin{bmatrix} x_0 & b\sigma(x_1) & d\tau(x_2) & bd\sigma\tau(x_{12}) \\ x_1 & \sigma(x_0) & d\tau(x_{12}) & d\sigma\tau(x_2) \\ x_2 & b\sigma(x_{12}) & \tau(x_0) & b\sigma\tau(x_1) \\ x_{12} & \sigma(x_2) & \tau(x_1) & \sigma\tau(x_0) \end{bmatrix} \right\},$$

where $x_0, x_1, x_2, x_{12} \in K$.

The proof is similar to the proof of Theorem 5.1 but this time the “hopping over” rules (5) should be used.

VI. SPRINGER’S THEOREM

So far we have described quaternion and biquaternion algebras over arbitrary fields (in which 2 is invertible) and shown how to compute their matrix representations. In addition we have seen that a given quaternion or biquaternion algebra is a *division algebra* if and only if its norm form or Albert form, respectively, is anisotropic.

The next question is how to determine when a quadratic form over a given field is actually anisotropic. In general this is a difficult problem, but for the fields in which we are interested we are in luck! The idea is to embed them into bigger fields that are complete with respect to a discrete valuation. For such fields there exists an easy criterion for anisotropy: Springer’s theorem. If the quadratic form is anisotropic over the bigger field, then it was already anisotropic over the smaller field. Applying this to a norm form or Albert form we get that if a quaternion or biquaternion algebra is a division algebra over the bigger field, then it was already division over the smaller field.

We will now present the details of this approach. Our fields of interest are:

- number fields F (i.e., finite extensions of the field $\mathbb{Q}$ of rational numbers);
- rational function field extensions of F of the form $F(x, y)$.

These are *discretely valued fields*. We embed them into bigger fields:

- $F \hookrightarrow F_{\mathfrak{p}}$: $\mathfrak{p}$ -adic field;
- $F(x, y) \hookrightarrow F((x))((y))$: iterated Laurent series field.

These bigger fields are *complete with respect to a discrete valuation* (or CDV fields for short). They are the completions of F and $F(x, y)$, respectively, for an appropriate discrete valuation. We refer to Appendix A for more background.

Theorem 6.1 (Springer’s theorem on CDV fields): Let F be a CDV field with residue field $\overline{F}$, uniformizer π and group of units $\mathcal{U}$. Assume that F is non-dyadic (i.e., $\text{char}(\overline{F}) \neq 2$). Let φ be a quadratic form over F . Write

$$\varphi = \varphi_1 \perp \langle \pi \rangle \varphi_2$$

where $\varphi_1 = \langle u_1, \dots, u_r \rangle$, $\varphi_2 = \langle u_{r+1}, \dots, u_n \rangle$ and $u_i \in \mathcal{U}$ for $1 \leq i \leq n$. Then φ is anisotropic over F if and only if the “first residue form” $\overline{\varphi}_1 = \langle \overline{u}_1, \dots, \overline{u}_r \rangle$ and the “second residue form” $\overline{\varphi}_2 = \langle \overline{u}_{r+1}, \dots, \overline{u}_n \rangle$ are anisotropic over $\overline{F}$.

We refer to [6, Chapter VI, §1] for more information on Springer’s theorem.

VII. 2×2 STBCs FROM QUATERNION ALGEBRAS

Using Springer's theorem we can construct infinite families of quaternion division algebras over number fields. They will give rise to 2×2 STBCs by means of the matrix representation in Theorem 5.1. We refer to Appendix B for more background on number fields.

Theorem 7.1: Let F be a number field, $\mathfrak{p}$ a prime ideal of F and π an element of F such that $v_{\mathfrak{p}}(\pi) = 1$. Then for any element $b \in F$ which is not a square modulo $\mathfrak{p}$, the quaternion algebra $(\pi, b)_F$ is a division algebra.

Proof: We show that the norm form $N = \langle 1, -\pi, -b, \pi b \rangle$ is anisotropic over F . Consider the $\mathfrak{p}$ -adic valuation on F . Let $\mathcal{O}_{\mathfrak{p}}$ denote the valuation ring. We obtain the following decomposition of N in $F_{\mathfrak{p}}$:

$$N = \varphi_1 \perp \langle \pi \rangle \varphi_2,$$

where $\varphi_1 = \langle 1, -b \rangle$ and $\varphi_2 = \langle -1, b \rangle$. Let $\overline{\varphi}_i$ denote the reduction of φ_i modulo $\mathfrak{p}$. Since we chose b to be a non-square in $\mathcal{O}_F/\mathfrak{p}$, the equation $x^2b = y^2$ has no nontrivial solutions in $\mathcal{O}_F/\mathfrak{p}$, therefore both $\overline{\varphi}_1$ and $\overline{\varphi}_2$ are anisotropic in $\mathcal{O}_F/\mathfrak{p}$. By Springer's theorem it follows that N is anisotropic over $F_{\mathfrak{p}}$. Since F is embedded in $F_{\mathfrak{p}}$, any nontrivial zero of N over F would give rise to a nontrivial zero over $F_{\mathfrak{p}}$. It follows that N is anisotropic over F . Hence $(\pi, b)_F$ is a division algebra. ■

Corollary 7.2: Let F be a number field. There are infinitely many quaternion division algebras $(\pi, b)_F$.

Proof: For any number field F , there are infinitely many primes $\mathfrak{p}$ in $\mathcal{O}_F$. Since any element of $\mathcal{O}_F$ is contained in only finitely many primes, this gives rise to infinitely many elements $\pi = \pi_{\mathfrak{p}}$. Having picked π , we can always find $b \in \mathcal{O}_F$ which is non-square modulo π . This is because the map $x \mapsto x^2$ in $\mathcal{O}_F/\mathfrak{p}$ is never surjective (since $\mathfrak{p}$ is not a prime above 2, it has a nontrivial kernel containing -1). ■

In the examples below we illustrate how to use Theorem 7.1. We start with the simplest case when the number field F is $\mathbb{Q}$.

Example 7.3: The quaternion algebra $(3, 5)_{\mathbb{Q}}$ is a division algebra.

Proof: The ideal $\mathfrak{p}$ in this case is simply $3\mathbb{Z}$, with uniformizer 3. Let $\overline{x}$ denote the reduction of x modulo $\mathfrak{p}$. The squares in $\mathbb{Z}/3\mathbb{Z}$ are $\{0, 1\}$. Since $5 \equiv 2 \pmod{3}$, it is not a square, hence the conditions of Theorem 7.1 are satisfied. We conclude that $(3, 5)_{\mathbb{Q}}$ is a division algebra. ■

Note that instead of 5 we could use any element congruent to 2 modulo 3.

In the next example we construct an infinite family of quaternion division algebras over $\mathbb{Q}(i)$, where $i = \sqrt{-1}$.

Example 7.4: Let $\mathfrak{p} = (2 + i)\mathbb{Z}[i]$ and let $b \equiv \pm i \pmod{\mathfrak{p}}$. Then $(2 + i, b)_{\mathbb{Q}(i)}$ is a division algebra.

Proof: Note that $2 + i$ is prime in $\mathbb{Z}[i]$ since its norm, 5, is prime in $\mathbb{Z}$. Hence $\mathfrak{p}$ is a prime ideal of $\mathbb{Z}[i]$. We need to show that $2 + i$ is not a square in $\mathbb{Z}[i]/\mathfrak{p}$. Note that $\mathfrak{p}$ lies above the prime ideal $5\mathbb{Z}$: by Theorem B.3 we have $5\mathbb{Z}[i] = (2 + i)(2 - i) = \mathfrak{p}\overline{\mathfrak{p}}$. In addition the inertial degree of $\mathfrak{p}$ over $5\mathbb{Z}$ is equal to 1. Hence the residue field $\mathbb{Z}[i]/\mathfrak{p}$ is a trivial extension of $\mathbb{Z}/5\mathbb{Z}$. Thus $\mathbb{Z}[i]/\mathfrak{p}$ has order 5 and can be represented by the elements $\{0, 1, -1, i, -i\}$. It follows that the squares in

$\mathbb{Z}[i]/(2 + i)$ are precisely $0, 1, -1$. By our choice of b and Theorem 7.1 we are done. ■

In the next example we pick a prime ideal $\mathfrak{p}$ of $\mathbb{Z}[i]$ with a nontrivial inertial degree in order to construct another infinite family of quaternion division algebras over $\mathbb{Q}(i)$.

Example 7.5: Let $\mathfrak{p} = 7\mathbb{Z}[i]$. There are 24 nonsquares in $\mathbb{Z}[i]/\mathfrak{p}$ and for any $b \in \mathbb{Z}[i]$ which is not a square modulo $\mathfrak{p}$, we have $(7, b)_{\mathbb{Q}(i)}$ is a division algebra.

Proof: By Theorem B.3, the ideal $\mathfrak{p} = 7\mathbb{Z}[i]$ is the principal ideal generated by 7 and its inertial degree over $7\mathbb{Z}$ is 2. Hence the number of elements in $\mathbb{Z}[i]/\mathfrak{p}$ is $7^2 = 49$. As an additive group the residue field $\mathbb{Z}[i]/\mathfrak{p}$ is isomorphic to $\mathbb{Z}/7\mathbb{Z} \oplus \mathbb{Z}/7\mathbb{Z}$: it has 49 elements of the form $\overline{a} + \overline{b}i$ with $\overline{a}, \overline{b} \in \mathbb{Z}/7\mathbb{Z}$. Consider the map $s : \mathbb{Z}[i]/\mathfrak{p}^{\times} \rightarrow \mathbb{Z}[i]/\mathfrak{p}^{\times}$ of the multiplicative group $\mathbb{Z}[i]/\mathfrak{p}^{\times}$, which sends every element to its square. The kernel of s is of order 2 and therefore the order of the image is $|\mathbb{Z}[i]/\mathfrak{p}^{\times}|/2 = 24$. More precisely, the image consists of 24 nonzero squares in $\mathbb{Z}[i]/\mathfrak{p}$, which all have the form $(\overline{x} + \overline{y}i)^2 = (\overline{x}^2 - \overline{y}^2) + i2\overline{x}\overline{y}$, where $x, y \in \mathbb{Z}$ and the bar denotes reduction modulo 7. This leaves 24 choices for the congruence class of b which satisfy the condition of Theorem 7.1. Note that picking an ideal $\mathfrak{p}$ with a nontrivial inertial degree gives rise to more candidates for the congruence class of b in $\mathbb{Z}[i]/\mathfrak{p}$. ■

VIII. 4×4 STBCs FROM BIQUATERNION ALGEBRAS

The following theorem can be used to construct 4×4 STBCs by means of the matrix representation in Theorem 5.1.

Theorem 8.1: Let $K = F(x, y)$, where F is a number field, and x, y are transcendental over F . Let $B = (a, x)_K \otimes (b, y)_K$. If $a, b \in F^{\times}$ are such that $a, b, ab \notin F^{\times 2}$, then B is a division algebra over K .

Proof: The biquaternion algebra B will be a division algebra over K if and only if its Albert form

$$\varphi = \langle a, x, -ax, -b, -y, by \rangle$$

is anisotropic over K . We will show that this is the case by an iterated application of Springer's Theorem.

Consider the embedding of $K = F(x, y)$ into the iterated Laurent series field $F((x))((y))$. (Note that $F((x))((y)) \neq F((y))((x))$.) Let v_y denote the y -adic valuation of $F((x))((y))$. We obtain the decomposition

$$\varphi = \varphi_1 \perp \langle y \rangle \varphi_2,$$

where $\varphi_1 = \langle a, x, -ax, -b \rangle$ and $\varphi_2 = \langle -1, b \rangle$. The form φ will be anisotropic over $F((x))((y))$ if and only if the forms $\overline{\varphi}_1$ and $\overline{\varphi}_2$ are anisotropic over $F((x))$, the residue field of v_y . The form $\overline{\varphi}_2$ is clearly anisotropic over $F((x))$ by our assumption. We can decompose the form $\overline{\varphi}_1$ as

$$\overline{\varphi}_1 = \psi_1 \perp \langle x \rangle \psi_2,$$

where $\psi_1 = \langle a, -b \rangle$ and $\psi_2 = \langle 1, -a \rangle$. By our assumption the residue forms $\overline{\psi}_1$ and $\overline{\psi}_2$ are clearly anisotropic over F so that $\overline{\varphi}_1$ is anisotropic over $F((x))$. In conclusion, the form φ is anisotropic over $F((x))((y))$, and thus certainly anisotropic over the smaller field K . ■

Remark 8.2: Note that picking transcendental elements of the form $x = e^{i\alpha}, y = e^{i\beta}$ guarantees that they have Euclidean norm 1. In the context of STBCs this can be used to satisfy uniform energy constraints.

APPENDIX A

PRELIMINARIES FROM DISCRETE VALUATION THEORY

We refer to [7, Chapter 1] for the details. Let F be a field.

A *discrete valuation* on F is a map

$$v : F \rightarrow \mathbb{Z} \cup \{\infty\}$$

with the following properties:

- $v(a) = \infty \Leftrightarrow a = 0$,
- $v(ab) = v(a) + v(b)$,
- $v(a + b) \geq \min(v(a), v(b))$.

Given a valuation v on F , let

$$\begin{aligned} \mathcal{O}_v &:= \{a \in F \mid v(a) \geq 0\}, \\ \mathfrak{m}_v &:= \{a \in F \mid v(a) > 0\}, \text{ and} \\ \mathcal{U}_v &:= \{a \in F \mid v(a) = 0\}. \end{aligned}$$

The objects just defined are the *valuation ring*, the *maximal ideal* and the *group of units* of v , respectively. The ring $\mathcal{O}_v$ is a local ring and $\mathfrak{m}_v$ is its unique maximal ideal. The *residue field* of v , $\overline{F}_v$, is defined to be $\mathcal{O}_v/\mathfrak{m}_v$.

Any discrete valuation v on F induces an *absolute value* $|a| := c^{v(a)}$, where $0 < c < 1$ is a real constant. Two absolute values $|\cdot|_1, |\cdot|_2$ are said to be equivalent if there is a constant t such that $|a|_1 = |a|_2^t$ for all $a \in F$. Let $|\cdot|_v$ denote the equivalence class of absolute values induced by v . If v is a discrete valuation on F , then F has a completion F_v with respect to $|\cdot|_v$. This completion is unique up to an isomorphism over F . An element π of F_v such that $v(\pi) = 1$ is called a *uniformizer*. Any two uniformizers differ by multiplication by a unit. Any element $x \in F_v^\times$ has the form $x = \pi^i u$, where i is an integer and $u \in \mathcal{U}_v$ is a unit. This gives us the following isomorphism of the multiplicative group of $F_v^\times$:

$$F_v^\times \cong \mathbb{Z} \times \mathcal{U}_v.$$

Below we give some examples of discrete valuations.

Example A.1: Consider a prime ideal $p\mathbb{Z}$ of $\mathbb{Z}$. Any nonzero rational number can be expressed as $p^a x/y$, where a, x and y are integers and $p \nmid xy$. Define the *p-adic valuation* on $\mathbb{Q}$ by $v_p : p^a x/y \mapsto a$. Then

$$\begin{aligned} \mathcal{O}_{v_p} &= \left\{ \frac{x}{y} \in \mathbb{Q} \mid p \nmid y \right\}, \\ \mathfrak{m}_{v_p} &= \left\{ \frac{x}{y} \in \mathbb{Q} \mid p \mid x \right\}, \text{ and} \\ \mathcal{U}_{v_p} &= \left\{ \frac{x}{y} \in \mathbb{Q} \mid p \nmid xy \right\}. \end{aligned}$$

$\mathbb{Q}_p$ denotes the completion of $\mathbb{Q}$ with respect to the p -adic valuation and $\mathbb{Z}_p$ denotes the ring of integers of $\mathbb{Q}_p$. The residue field $\overline{\mathbb{Q}}_p \cong \mathbb{Z}_p/p\mathbb{Z}_p \cong \mathbb{F}_p$ is a finite field of order p .

Similarly we can define *p-adic valuations* on number fields, as in the next example.

Example A.2: Let F be a number field with ring of integers $\mathcal{O}_F$. Any prime ideal $\mathfrak{p}$ of $\mathcal{O}_F$ induces a discrete valuation on F . Namely, let $v_{\mathfrak{p}}$ map a nonzero element x of $\mathcal{O}_F$ to the largest integer i , such that $x \in \mathfrak{p}^i$, and let $v_{\mathfrak{p}}(0) = \infty$. Extend this map to F by letting $v_{\mathfrak{p}}(x/y) := v_{\mathfrak{p}}(x) - v_{\mathfrak{p}}(y)$. The map just defined is a discrete valuation on F . We let $F_{\mathfrak{p}}$ denote $F_{v_{\mathfrak{p}}}$, the completion of F under $v_{\mathfrak{p}}$. Similarly let $\mathcal{O}_{\mathfrak{p}}$ and $\mathcal{U}_{\mathfrak{p}}$ denote the ring of integers of $F_{\mathfrak{p}}$ (also called *p-adic integers*) and the group of units of $\mathcal{O}_{\mathfrak{p}}$ (called *p-adic units*), respectively.

Example A.3: Let F be a field and let $K = F(x)$. For

$$K \ni f(x) = \sum_{i=m}^N a_i x^i, \quad a_m \neq 0, m \in \mathbb{Z}$$

define $v_x(f) = m$. Then $\mathcal{O}_{v_x} = F[x]$, with residue field F . The completion of K with respect to v_x is the field of Laurent series, denoted by $F((x))$. Its elements are of the form $\sum_{i=m}^{\infty} a_i x^i$ with $m \in \mathbb{Z}$. The valuation ring of $F((x))$ with respect to v_x is the ring of formal power series $F[[x]]$, whose elements are of the form $\sum_{i=0}^{\infty} a_i x^i$. Its residue field is F .

APPENDIX B

PRELIMINARIES FROM NUMBER THEORY

Next we look at how prime ideals behave in extensions of number fields. Let F be a number field, i.e., a finite extension of $\mathbb{Q}$. Let K be a finite extension of F . If $\mathfrak{p}$ is a prime ideal of $\mathcal{O}_F$, then $\mathfrak{p}\mathcal{O}_K$ is an ideal of $\mathcal{O}_K$ with unique factorization into prime ideals of $\mathcal{O}_K$:

$$\mathfrak{p}\mathcal{O}_K = \mathfrak{P}_1^{e_1} \cdots \mathfrak{P}_r^{e_r}.$$

The integer e_i is called the *ramification index* of $\mathfrak{P}_i$ over $\mathfrak{p}$. The *inertial degree* f_i is defined to be $[\mathcal{O}_K/\mathfrak{P}_i : \mathcal{O}_F/\mathfrak{p}]$, the degree of the field extension $\mathcal{O}_K/\mathfrak{P}_i$ of $\mathcal{O}_F/\mathfrak{p}$.

When K/F is a Galois extension, the ramification indices (resp. inertial degrees) are the same for all i and we denote them simply by $e_{\mathfrak{p}}$ (resp. $f_{\mathfrak{p}}$). The following theorem describes the relationship between inertial and ramification degrees in this case.

Theorem B.1 ([8, Theorem 21]): Let K be a Galois extension of F of degree n and let $\mathfrak{p}$ be a prime ideal of F . Then $\mathfrak{p}\mathcal{O}_K = (\mathfrak{P}_1 \cdots \mathfrak{P}_{r_{\mathfrak{p}}})^{e_{\mathfrak{p}}}$ with $r_{\mathfrak{p}} e_{\mathfrak{p}} f_{\mathfrak{p}} = n = [K : F]$.

When the prime ideal in the ground field F is understood, we may drop the indices and refer to these parameters as r, e, f .

In the special case of cyclotomic extensions of $\mathbb{Q}$ we have the following result:

Theorem B.2 ([8, Theorem 26]): Let $F = \mathbb{Q}(\zeta_m)$, where ζ_m is a primitive m -th root of unity. Let p be a prime not dividing m . Then the inertial degree f_p in F is the multiplicative order of p modulo m .

Finally, for the decomposition of primes in the extension $\mathbb{Q}(i)$ of $\mathbb{Q}$ (with $i^2 = -1$) we obtain the following theorem as a special case of [8, Theorem 25]:

Theorem B.3: Let $F = \mathbb{Q}(i)$ (and thus $\mathcal{O}_F = \mathbb{Z}[i]$) and let p be an odd prime.

If $p \equiv 1 \pmod{4}$ then

$$p\mathbb{Z}[i] = (\pi)(\bar{\pi}) = \mathfrak{p}\bar{\mathfrak{p}},$$

where π and $\bar{\pi}$ are non-associated, prime and of norm p in $\mathbb{Z}[i]$. Thus $r_p = 2$, $f_p = e_p = 1$.

If $p \equiv 3 \pmod{4}$ then

$$p\mathbb{Z}[i] = (p),$$

i.e., the ideal $p\mathbb{Z}$ remains inert in $\mathbb{Z}[i]$ and $r_p = e_p = 1$, $f_p = 2$.

REFERENCES

- [1] S.M. Alamouti, "A simple transmit diversity technique for wireless communications," *IEEE J. Sel. Areas Commun.*, vol. 16, no. 8, pp. 1451–1458, 1998.
- [2] B.A. Sethuraman, B.S. Rajan and V. Shashidhar, "Full-diversity, high-rate space-time block codes from division algebras," *IEEE Trans. Inf. Theory*, vol. 49, no. 10, pp. 2596–2616, 2003, Special Issue on Space-Time Transmission, Reception, Coding and Signal Design.
- [3] V. Shashidhar, B.S. Rajan and B.A. Sethuraman, "Information-lossless space-time block codes from crossed-product algebras," *IEEE Trans. Inf. Theory*, vol. 52, no. 9, pp. 3913–3935, 2006.
- [4] G. Berhuy and F.E. Oggier, "Space-Time Codes from Crossed Product Algebras of Degree 4," *Applied Algebra, Algebraic Algorithms and Error-Correcting Codes*, pp. 90–99, 2007.
- [5] D.W. Lewis, "Quaternion algebras and the algebraic legacy of Hamilton's quaternions," *Irish Math. Soc. Bull.*, no. 57, pp. 41–64, 2006.
- [6] T.Y. Lam, *Introduction to Quadratic Forms over Fields*. Providence, RI: American Mathematical Society, 2005, Graduate Studies in Mathematics no. 67.
- [7] I.B. Fesenko and S.V. Vostokov, *Local Fields and Their Extensions*. With a foreword by I. R. Shafarevich. Second edition. Providence, RI: American Mathematical Society, 2002, Translations of Mathematical Monographs no. 121.
- [8] D.A. Marcus, *Number Fields*. New York-Heidelberg: Springer-Verlag, 1977, Universitext.